

Where to look for new physics?

Indications from Dark Matter and extra Higgses

Filippo Sala

CNRS and IPhT, CEA/Saclay



Fermilab, 3 Sept 2015



The Standard Model



?The Standard Model?

? Supersymmetry ? ??? ? Composite models ?

Any indication of a New Physics scale?



Data: no answer, but something has to be there, like Dark Matter

Theory: answer “ \equiv ” attitude towards the hierarchy problem

Any indication of a New Physics scale?



Data: no answer, but something has to be there, like Dark Matter

Theory: answer “ \equiv ” attitude towards the hierarchy problem

$$M_{\text{NP}} = 0.1 \div 10 \text{ TeV}$$

Protect the mass of the scalars from any NP [$'t$ Hooft 1979, ...]

The Fermi scale is **natural** Fine-tuning Δ “small” $\Rightarrow M_{\text{NP}}$ “small”

Examples Supersymmetry Twin and composite Higgs models

No evidence for NP puts pressure on this attitude

Any indication of a New Physics scale?



Data: no answer, but something has to be there, like Dark Matter

Theory: answer “ \equiv ” attitude towards the hierarchy problem

$$M_{\text{NP}} = 0.1 \div 10 \text{ TeV}$$

Protect the mass of the scalars from any NP [*'t Hooft 1979, ...*]

The Fermi scale is **natural** Fine-tuning Δ “small” $\Rightarrow M_{\text{NP}}$ “small”

Examples Supersymmetry Twin and composite Higgs models

No evidence for NP puts pressure on this attitude


$$M_{\text{NP}} \text{ can be } \gg \text{TeV}$$

- Think different (e.g. “UV naturalness”, cosmological relaxation)
- Accept the tuning Δ (and go anthropic)


Future lampposts in this talk



Colliders



Telescopes



Where to look for New Physics?

→ Indications from “natural” New Physics

mostly based on Buttazzo S Tesi, 1505.05488

→ Indications from WIMP Dark Matter

mostly based on Cirelli Hambye Panci S Taoso, 1507.05519

Where to look for New Physics?

→ Indications from “natural” New Physics

mostly based on Buttazzo S Tesi, 1505.05488

- ◊ Can new Higgses be the lightest new particles around?
- ◊ How to look for them?

- ◊ Can new Higgses be the lightest new particles around?

Another light scalar would be hard to justify with anthropic arguments

- ◊ Can new Higgses be the lightest new particles around?

Another light scalar would be hard to justify with anthropic arguments

Extra **Singlet**-like Higgses are ubiquitous, for example in

- Twin Higgs
- Supersymmetry
- Electroweak Baryogenesis (independent of naturalness)

Why interesting? Solves little hierarchy, without coloured top partners

If nothing new at the LHC14, TH models still quite natural!

Why interesting? Solves little hierarchy, without coloured top partners

If nothing new at the LHC14, TH models still quite natural!

→ Add a Z_2 -symmetric copy of the SM

[only copy of top strictly necessary see e.g. J Serra @ MIAPP 2015]

Why interesting? Solves little hierarchy, without coloured top partners

If nothing new at the LHC14, TH models still quite natural!

- Add a Z_2 -symmetric copy of the SM
 - [only copy of top strictly necessary see e.g. J Serra @ MIAPP 2015]
- 8 “Higgs” degrees of freedom - vs 4 in the SM
 - 7 are massless Goldstone bosons
 - one, $\sigma =$ radial mode of $\mathcal{G} \rightarrow \mathcal{H}$
 - $\langle \sigma \rangle = f$, $m_\sigma \sim f$ conceivable if UV completion is weakly coupled

Why interesting? Solves little hierarchy, without coloured top partners

If nothing new at the LHC14, TH models still quite natural!

- Add a Z_2 -symmetric copy of the SM
 - [only copy of top strictly necessary see e.g. J Serra @ MIAPP 2015]
- 8 “Higgs” degrees of freedom - vs 4 in the SM
 - 7 are massless Goldstone bosons
 - one, $\sigma =$ radial mode of $\mathcal{G} \rightarrow \mathcal{H}$
 - $\langle \sigma \rangle = f$, $m_\sigma \sim f$ conceivable if UV completion is weakly coupled
- Small explicit breaking of both \mathcal{G} and Z_2 to get m_h and $v_{EW} \neq 0$

Why interesting? Solves little hierarchy, without coloured top partners

If nothing new at the LHC14, TH models still quite natural!

- Add a Z_2 -symmetric copy of the SM
 - [only copy of top strictly necessary see e.g. J Serra @ MIAPP 2015]
- 8 “Higgs” degrees of freedom - vs 4 in the SM
 - 7 are massless Goldstone bosons
 - one, $\sigma =$ radial mode of $\mathcal{G} \rightarrow \mathcal{H}$
 - $\langle \sigma \rangle = f$, $m_\sigma \sim f$ conceivable if UV completion is weakly coupled
- Small explicit breaking of both \mathcal{G} and Z_2 to get m_h and $v_{EW} \neq 0$

Other particles? Either $M \gtrsim 4\pi f$ or very weakly coupled

Light extra Higgses in Supersymmetry

MSSM

$$m_h^2 \leq m_Z^2 \cos^2 2\beta + \Delta_t^2 \Rightarrow \Delta_t \gtrsim 85 \text{ GeV} \Rightarrow \text{stops heavier than } \sim 1.5 \text{ TeV}$$

Fine tuning worse than 1%!

$$\frac{dv^2}{dm_{H_u}^2} \simeq \frac{4}{g^2}$$

$$\delta m_{H_u}^2 \sim -\frac{3y_t^2}{4\pi^2} \cancel{m_t^2} \log \frac{\Lambda}{m_{\tilde{t}}}$$

Light extra Higgses in Supersymmetry

MSSM

$$m_h^2 \leq m_Z^2 \cos^2 2\beta + \Delta_t^2 \Rightarrow \Delta_t \gtrsim 85 \text{ GeV} \Rightarrow \text{stops heavier than } \sim 1.5 \text{ TeV}$$

Fine tuning worse than 1%!

$$\frac{dv^2}{dm_{H_u}^2} \simeq \frac{4}{g^2}$$

$$\delta m_{H_u}^2 \sim -\frac{3y_t^2}{4\pi^2} \cancel{m_t^2} \log \frac{\Lambda}{m_t}$$


NMSSM

Add a singlet $S \quad \Delta W = \lambda S H_u H_d + f(S)$

$$m_h^2 \leq m_Z^2 c_{2\beta}^2 + \Delta_t^2 + \cancel{\lambda}^2 v^2 s_{2\beta}^2$$

Fine tuning better than 5%!
[green points, $\tan \beta \lesssim 5$]

$$\frac{dv^2}{dm_{H_u}^2} \simeq \frac{\kappa}{\cancel{\lambda}^3} \frac{1}{t_{2\beta}}$$




Gherghetta et al. 1212.5243 [$\Lambda = 20$ TeV]

NMSSM spectrum

NMSSM with $\lambda \sim 1$ and heavy stops & gluinos

[$\lambda \gtrsim 0.7$ needs a completion before GUT scale!]



NMSSM spectrum

NMSSM with $\lambda \sim 1$ and heavy stops & gluinos


[$\lambda \gtrsim 0.7$ needs a completion before GUT scale!]

The scalars are:

CP-even h, h_3, ϕ (from h_ν, H, S)

CP-odd A, A_s

H^\pm



NMSSM spectrum


NMSSM with $\lambda \sim 1$ and heavy stops & gluinos

[$\lambda \gtrsim 0.7$ needs a completion before GUT scale!]

The scalars are:

CP-even h, h_3, ϕ (from h_ν, H, S)

$$\mathcal{H}_{\text{ph}} \equiv \begin{pmatrix} h_3 \\ h \\ \phi \end{pmatrix} = R^T \begin{pmatrix} H \\ h_\nu \\ S \end{pmatrix}, \quad R = [R_\delta^{12} R_\gamma^{23} R_\sigma^{13}]$$



NMSSM spectrum


NMSSM with $\lambda \sim 1$ and heavy stops & gluinos

[$\lambda \gtrsim 0.7$ needs a completion before GUT scale!]


The scalars are:

CP-even h, h_3, ϕ (from h_ν, H, S)

$$\mathcal{H}_{\text{ph}} \equiv \begin{pmatrix} h_3 \\ h \\ \phi \end{pmatrix} = R^T \begin{pmatrix} H \\ h_\nu \\ S \end{pmatrix}, \quad R = \boxed{R_\delta^{12} R_\gamma^{23} R_\sigma^{13}}$$



$\gamma = h_\nu - S$ mixing



A motivated limiting case

$m_{h_3} \gg m_{h,\phi}$ and $\sigma, \delta \rightarrow 0$


Bottom-up motivation for a Singlet: Higgs couplings fit

Bottom-up motivation for a Singlet: Higgs couplings fit


$$h_{\text{LHC}} = h = c_\gamma(c_\delta h_\nu - s_\delta H) + s_\gamma S$$

$$\left[\frac{g_{htt}}{g_{htt}^{\text{SM}}} = c_\gamma(c_\delta + \frac{s_\delta}{\tan \beta}), \quad \frac{g_{hbb}}{g_{hbb}^{\text{SM}}} = c_\gamma(c_\delta - s_\delta \tan \beta), \quad \frac{g_{hVV}}{g_{hVV}^{\text{SM}}} = c_\gamma c_\delta \right]$$

LHC8 status



$$s_\gamma^2 = 0, 0.15, 0.3$$





Bottom-up motivation for a Singlet: Higgs couplings fit

$$h_{\text{LHC}} = h = c_\gamma(c_\delta h_\nu - s_\delta H) + s_\gamma S$$

$$\left[\frac{g_{htt}}{g_{SM}^{tt}} = c_\gamma(c_\delta + \frac{s_\delta}{\tan \beta}), \quad \frac{g_{hbb}}{g_{SM}^{bb}} = c_\gamma(c_\delta - s_\delta \tan \beta), \quad \frac{g_{hVV}}{g_{SM}^{VV}} = c_\gamma c_\delta \right]$$

LHC14 projections (300 fb^{-1})




- ✓ Can new Higgses be the lightest new particles around?


- ◊ How to look for them?

- ◊ How to look for them?

At which machines?



Currently unclear where particle physicists will put (EU? China? ???) money:



HL-LHC $\sqrt{s} = 14 \text{ TeV}$, 3000 fb^{-1} , $\sim 2025\text{-}2035$

HE-LHC $\sqrt{s} = 33 \text{ TeV}$, needs new technology

FCC-hh $\sqrt{s} \sim 100 \text{ TeV}$, start $\sim 2040(?)$,
needs $\sim 100 \text{ km}$ tunnel

ILC $\sqrt{s} = 0.5 - 1 \text{ TeV}$, maybe Japan soon

CLIC \sqrt{s} up to 3 TeV, needs new technology

FCC-ee \sqrt{s} up to 500 GeV, higher luminosity,
needs $\sim 100 \text{ km}$ tunnel

Generic singlet

$$\sin^2 \gamma = \frac{M_{hh}^2 - m_\phi}{m_\phi^2 - m_h^2}$$

Master formula, valid for **any** model

2 free parameters control all pheno! + $\text{BR}_{\phi \rightarrow hh}$ ($= \text{BR}_{\phi \rightarrow ZZ}$ at $m_\phi \gg m_W$)

Generic singlet

$$\sin^2 \gamma = \frac{M_{hh}^2 - m_\phi}{m_\phi^2 - m_h^2}$$

Master formula, valid for **any** model

2 free parameters control all pheno! + $\text{BR}_{\phi \rightarrow hh}$ ($= \text{BR}_{\phi \rightarrow ZZ}$ at $m_\phi \gg m_W$)

h : signal strengths $\mu = c_\gamma^2 \times \mu_{\text{SM}}$ ϕ : $\mu(m_\phi) = s_\gamma^2 \times \mu_{\text{SM}}(m_\phi)$ [barring $\phi \rightarrow hh$]

Generic singlet

$$\sin^2 \gamma = \frac{M_{hh}^2 - m_\phi^2}{m_\phi^2 - m_h^2}$$

Master formula, valid for **any** model

2 free parameters control all pheno! + $\text{BR}_{\phi \rightarrow hh}$ ($= \text{BR}_{\phi \rightarrow ZZ}$ at $m_\phi \gg m_W$)

h : signal strengths $\mu = c_\gamma^2 \times \mu_{\text{SM}}$ ϕ : $\mu(m_\phi) = s_\gamma^2 \times \mu_{\text{SM}}(m_\phi)$ [barring $\phi \rightarrow hh$]

What does one learn from the potential $f(S)$?

$$\text{BR}_{\phi \rightarrow hh} = \frac{1}{4} - \frac{3}{4} \frac{\nu}{v_s} \frac{\sqrt{M_{hh}^2 - m_h^2}}{m_\phi} + O\left(\frac{\nu^2}{m_\phi^2}\right)$$

$$\frac{g_{h^3}}{g_{h^3}^{\text{SM}}} = 1 + \frac{2}{3} \frac{\nu}{v_s} \frac{\sqrt{M_{hh}^2 - m_h^2}}{m_\phi} \left(\frac{M_{hh}^2}{m_h^2} - 1 \right) + O\left(\frac{\nu^2}{m_\phi^2}\right)$$

Valid for **any** potential!! v_s leading new parameter

Generic singlet: Higgs couplings

$$\sin^2 \gamma = \frac{M_{hh}^2 - m_h^2}{m_\phi^2 - m_h^2}$$

Master formula, valid for **any** model

2 free parameters control all pheno! + $\text{BR}_{\phi \rightarrow hh}$ ($= \text{BR}_{\phi \rightarrow ZZ}$ at $m_\phi \gg m_W$)

h: signal strengths $\mu = c_\gamma^2 \times \mu_{\text{SM}}$



Generic singlet: Higgs couplings

$$\sin^2 \gamma = \frac{M_{hh}^2 - m_h^2}{m_\phi^2 - m_h^2}$$

Master formula, valid for **any** model

2 free parameters control all pheno! + $\text{BR}_{\phi \rightarrow hh}$ ($= \text{BR}_{\phi \rightarrow ZZ}$ at $m_\phi \gg m_W$)

h : signal strengths $\mu = c_\gamma^2 \times \mu_{\text{SM}}$



Generic singlet: Higgs couplings


$$\sin^2 \gamma = \frac{M_{hh}^2 - m_h^2}{m_\phi^2 - m_h^2}$$

Master formula, valid for **any** model


2 free parameters control all pheno! + $\text{BR}_{\phi \rightarrow hh}$ ($= \text{BR}_{\phi \rightarrow ZZ}$ at $m_\phi \gg m_W$)

h : signal strengths $\mu = c_\gamma^2 \times \mu_{\text{SM}}$

Add g_{hhh} : could be first deviation seen!



Snowmass 2013



Generic singlet: Higgs couplings



$$\sin^2 \gamma = \frac{M_{hh}^2 - m_h^2}{m_\phi^2 - m_h^2}$$

Master formula, valid for **any** model

2 free parameters control all pheno! + $\text{BR}_{\phi \rightarrow hh}$ ($= \text{BR}_{\phi \rightarrow ZZ}$ at $m_\phi \gg m_W$)

h : signal strengths $\mu = c_\gamma^2 \times \mu_{\text{SM}}$

Add g_{hhh} : could be first deviation seen!



Generic singlet: direct searches

$$\sin^2 \gamma = \frac{M_{hh}^2 - m_h^2}{m_\phi^2 - m_h^2}$$

Master formula, valid for **any** model

2 free parameters control all pheno! + $\text{BR}_{\phi \rightarrow hh}$ ($= \text{BR}_{\phi \rightarrow ZZ}$ at $m_\phi \gg m_W$)

ϕ : $\mu(m_\phi) = s_\gamma^2 \times \mu_{\text{SM}}(m_\phi)$ [barring $\phi \rightarrow hh$]

Generic singlet: direct searches



$$\sin^2 \gamma = \frac{M_{hh}^2 - m_h^2}{m_\phi^2 - m_h^2}$$

Master formula, valid for **any** model

2 free parameters control all pheno! + $\text{BR}_{\phi \rightarrow hh}$ ($= \text{BR}_{\phi \rightarrow ZZ}$ at $m_\phi \gg m_W$)

Large $\text{BR}_{\phi \rightarrow hh}$ easier for $v_s < 0$

ϕ : $\mu(m_\phi) = s_\gamma^2 \times \mu_{\text{SM}}(m_\phi)$ [barring $\phi \rightarrow hh$]



Generic singlet: direct searches



$$\sin^2 \gamma = \frac{M_{hh}^2 - m_h^2}{m_\phi^2 - m_h^2}$$

Master formula, valid for **any** model

2 free parameters control all pheno! + $\text{BR}_{\phi \rightarrow hh}$ ($= \text{BR}_{\phi \rightarrow ZZ}$ at $m_\phi \gg m_W$)

Large $\text{BR}_{\phi \rightarrow hh}$ easier for $v_s < 0$

ϕ : $\mu(m_\phi) = s_\gamma^2 \times \mu_{\text{SM}}(m_\phi)$ [barring $\phi \rightarrow hh$]



Generic singlet: direct searches



$$\sin^2 \gamma = \frac{M_{hh}^2 - m_h^2}{m_\phi^2 - m_h^2}$$

Master formula, valid for **any** model

2 free parameters control all pheno! + $\text{BR}_{\phi \rightarrow hh}$ ($= \text{BR}_{\phi \rightarrow ZZ}$ at $m_\phi \gg m_W$)

h : signal strengths $\mu = c_\gamma^2 \times \mu_{\text{SM}}$

ϕ : $\mu(m_\phi) = s_\gamma^2 \times \mu_{\text{SM}}(m_\phi)$ [barring $\phi \rightarrow hh$]



Higgs as a PNG boson: Twin Higgs

Higgs as a PNG boson: Twin Higgs

Take-home message:

Mixing angle $\sin \gamma \simeq \frac{v}{f}$



Direct searches of the singlet are promising!

Higgs as a PNG boson: Twin Higgs

$$\sin^2 \gamma = \frac{M_{hh}^2 - m_h^2}{m_\sigma^2 - m_h^2}$$

$$M_{hh}^2 = (m_h^2 + m_\sigma^2) v^2/f^2$$

Only two free parameters f and m_σ \Rightarrow $\text{BR}_{\sigma \rightarrow hh}$ fixed everywhere

Twin SM \Rightarrow $\text{BR}_{\sigma \rightarrow \text{inv.}} \neq 0$ [equivalence theorem: $\text{BR}_{\sigma \rightarrow \text{inv.}} \rightarrow 3/7$ for $m_\sigma > m_Z \times f/v$]



Higgs as a PNG boson: Twin Higgs

$$\sin^2 \gamma = \frac{M_{hh}^2 - m_h^2}{m_\sigma^2 - m_h^2}$$

$$M_{hh}^2 = (m_h^2 + m_\sigma^2) v^2/f^2$$

Only two free parameters f and m_σ \Rightarrow $\text{BR}_{\sigma \rightarrow hh}$ fixed everywhere

Twin SM \Rightarrow $\text{BR}_{\sigma \rightarrow \text{inv.}} \neq 0$ [equivalence theorem: $\text{BR}_{\sigma \rightarrow \text{inv.}} \rightarrow 3/7$ for $m_\sigma > m_Z \times f/v$]



The NMSSM

$$\sin^2 \gamma = \frac{M_{hh}^2 - m_h^2}{m_\phi^2 - m_h^2}$$

$$M_{hh}^2 = m_Z^2 c_{2\beta}^2 + \lambda^2 v^2 s_{2\beta}^2 + \Delta^2$$

Remember: naturalness $\Rightarrow \lambda \sim 1$

Δ = all loop effects, e.g. top-stop

The NMSSM

$$\sin^2 \gamma = \frac{M_{hh}^2 - m_h^2}{m_\phi^2 - m_h^2}$$



$$M_{hh}^2 = m_Z^2 c_{2\beta}^2 + \lambda^2 v^2 s_{2\beta}^2 + \Delta^2$$

Remember: naturalness $\Rightarrow \lambda \sim 1$

Δ = all loop effects, e.g. top-stop

Here $\lambda = 1.2$ $\Delta = 70$ GeV

$\tan \beta$ “small” otherwise EWPT



The NMSSM

$$\sin^2 \gamma = \frac{M_{hh}^2 - m_h^2}{m_\phi^2 - m_h^2}$$



$$M_{hh}^2 = m_Z^2 c_{2\beta}^2 + \lambda^2 v^2 s_{2\beta}^2 + \Delta^2$$

Remember: naturalness $\Rightarrow \lambda \sim 1$

Δ = all loop effects, e.g. top-stop

Here $\lambda = 0.7$ $\Delta = 80$ GeV

$\tan \beta$ “small” otherwise EWPT




Fully mixed case and a $\gamma\gamma$ signal

Singlet-like state lighter than 125 GeV

Hard to see, but maybe already found??

SUSY 2015, Tristan du Pree






Fully mixed case and a $\gamma\gamma$ signal

Barbieri et al 1304.3670, 1307.4937

Singlet-like state lighter than 125 GeV

Hard to see, but maybe already found??

SUSY 2015, Tristan du Pree



Where to look for New Physics?

- ✓ Indications from “natural” New Physics

mostly based on Buttazzo S Tesi, 1505.05488


- Indications from WIMP Dark Matter

mostly based on Cirelli Hambye Panci S Taoso, 1507.05519

Where to look for New Physics?


- Indications from WIMP Dark Matter
mostly based on Cirelli Hambye Panci S Taoso, 1507.05519

Where is Dark Matter?




[courtesy of Marco Cirelli]

Where is Dark Matter?



[courtesy of Marco Cirelli]



How to probe the “thermal relic WIMP” paradigm?

[Unitarity bound: $M_{DM} < 80 \div 120$ TeV Griest Kamionkowski 1990,


...

Cahill-Rowley et al. 1501.03153]

General strategy: effective field theories?

The EFT approach:

- ⌚ Model-independent
- ⌚ easy comparison collider - direct detection




General strategy: effective field theories?

The EFT approach:

- ☺ Model-independent
- ☺ easy comparison collider - direct detection
- ☹ ~ wrong for LHC (especially 14 TeV) !!

often momentum transfer > suppression scale Λ



Lot of recent activity

- Busoni et al 1307.2253 and 1402.1275,
- Buchmuller et al 1308.6799,...
- Abdallah et al 1409.2893,
- Racco Wulzer Zwirner 1502.04701

Need to go to benchmark/simplified models!

Quantum numbers		
SU(2) _L	U(1) _Y	Spin
3	0	F
5	0	F

An EW fermion multiplet

Possibly the “simplest” simplified model

Philosophy: Focus on DM, and try to preserve SM successes (flavour & CP, ..)
+ DM stability, adding the least possible ingredients to the theory

Approach: add to the SM extra particle χ
and determine its “good” quantum numbers

“good” = i) stable ii) lightest component neutral iii) allowed

$$\mathcal{L} = \mathcal{L}_{\text{SM}} + c \bar{\chi}(i\hat{D} - M_\chi)\chi$$

other terms forbidden without new symmetries

M_χ is the only one free parameter, fixed if we impose thermal relic abundance!

$$M_{\text{thermal}}^{\text{5plet}} \simeq 9.4 \text{ TeV}$$

$$M_{\text{thermal}}^{\text{3plet}} \simeq 3 \text{ TeV}$$

Electroweak multiplets at colliders

5plet No hopes to reach M_{thermal}

3plet No hopes to reach M_{thermal} before a 100 TeV collider (i.e. before 2040)


Electroweak multiplets at colliders

5plet No hopes to reach M_{thermal}

3plet No hopes to reach M_{thermal} before a 100 TeV collider (i.e. before 2040)


Most promising channel: disappearing tracks

current stronger probe $M_\chi \gtrsim 270 \text{ GeV}$




Cirelli S Taoso 1407.7058

see also Feng Strassler 1994 ... Low Wang 1404.0682



Ostdiek 1506.03445

Direct Detection



Hisano et al. 1504.00915:

$$\sigma_{\text{SI}}^{\text{5plet}} = 1.9 \times 10^{-46} \text{ cm}^2$$

$$\sigma_{\text{SI}}^{\text{3plet}} = 2.3 \times 10^{-47} \text{ cm}^2$$

full NLO in α_S , $O(50\%)$ uncertainties [largest error from charm content of nucleon]

Electroweak multiplets in the (γ) sky





Sommerfeld enhancement

at low velocities non-rel. attractive potential

Milky Way $v \sim 10^{-3}c$

Dwarf spheroidals $v \sim 1 \div 5 \times 10^{-5}c$

$\chi_0\chi_0 \rightarrow WW, \gamma\gamma$ σv saturates at $v \lesssim 10^{-2}$ \rightarrow



Electroweak multiplets in the (γ) sky



Sommerfeld enhancement

at low velocities non-rel. attractive potential


Milky Way $v \sim 10^{-3}c$

Dwarf spheroidals $v \sim 1 \div 5 \times 10^{-5}c$


$\chi_0\chi_0 \rightarrow WW, \gamma\gamma$ σv saturates at $v \lesssim 10^{-2}$ \rightarrow



$\bar{p}, e^+, \nu, \gamma, \dots$



γ ray lines: smaller cross-sections





but features in γ spectrum enhance sensitivities

γ continuum with FERMI

- FERMI measures γ flux from all sky
- We carefully model astrophysical backgrounds (conservatively)
- We divide the sky into regions, and extract bounds from each one



γ continuum with FERMI

- FERMI measures γ flux from all sky
- We carefully model astrophysical backgrounds (conservatively)
- We divide the sky into regions, and extract bounds from each one



γ continuum with FERMI

- FERMI measures γ flux from all sky
- We carefully model astrophysical backgrounds (conservatively)
- We divide the sky into regions, and extract bounds from each one



- ◊ Galactic bounds depend on DM profile
- ◊ All bounds assume 5plet = 100% of DM

γ continuum from dwarf spheroidal galaxies

A primer on dwarf spheroidal galaxies

- ◊ gravitationally linked to our galaxy
- ◊ DM dominated objects → this is why they are good targets!
- ◊ often “trackers” are just a few → big uncertainties on DM properties


[with respect to Milky Way: almost no bkg, large uncertainties in J factors]

γ continuum from dwarf spheroidal galaxies

A primer on dwarf spheroidal galaxies

- ◊ gravitationally linked to our galaxy
- ◊ DM dominated objects → this is why they are good targets!
- ◊ often “trackers” are just a few → big uncertainties on DM properties

[with respect to Milky Way: almost no bkg, large uncertainties in J factors]



FERMI: 15 dwarves, assumes $\Delta J < 40\%$

HESS: subset of 4, plus Sagittarius


MAGIC: only Segue1 (large uncertainties!)

γ continuum from dwarf spheroidal galaxies

A primer on dwarf spheroidal galaxies

- ◊ gravitationally linked to our galaxy
- ◊ DM dominated objects → this is why they are good targets!
- ◊ often “trackers” are just a few → big uncertainties on DM properties


[with respect to Milky Way: almost no bkg, large uncertainties in J factors]





FERMI: 15 dwarves, assumes $\Delta J < 40\%$

HESS: subset of 4, plus Sagittarius

MAGIC: only Segue1 (large uncertainties!)



γ lines: galactic center and dwarves





[CTA prospects from [Ovanesyan et al 1409.8294](#) and [Bergstrom et al 1207.6773](#)]

MAGIC = only one that looked for lines from dwarves - but just Segue1

Lot of progress conceivable with dwarf spheroidals!

- Look at the same (other) dwarves with other (the same) experiments
- measure better DM properties to reduce uncertainties

γ lines: galactic center and dwarves - 3plet




[CTA prospects from [Ovanesyan et al 1409.8294](#) and [Bergstrom et al 1207.6773](#)]

MAGIC = only one that looked for lines from dwarves - but just Segue1

Lot of progress conceivable with dwarf spheroidals!

- Look at the same (other) dwarves with other (the same) experiments
- measure better DM properties to reduce uncertainties

An EW fermion 5plet: pheno summary



Where to look for New Physics?

- ✓ Indications from “natural” New Physics

Buttazzo S Tesi, 1505.05488

- ✓ Indications from WIMP Dark Matter

Cirelli Hambye Panci S Taoso, 1507.05519


Where to look for New Physics?

- ✓ Indications from “natural” New Physics
Buttazzo S Tesi, 1505.05488

An extra Singlet-like Higgs

appears in NMSSM, Twin Higgs, ...

$$h_2 \rightarrow VV, hh, \quad h \text{ couplings}, \quad g_{hhh}$$




- ✓ Indications from WIMP Dark Matter
Cirelli Hambye Panci S Taoso, 1507.05519

An EW fermion multiplet, focus on 5plet

- ◊ “simplest” simplified model?
- ◊ Minimal Dark Matter ◊ 3plet = Wino ...

If 100% of DM, dwarves might be the future



Back up Dark Matter

Minimal Dark Matter: candidates


Allowed: χ neutral under g, γ , and almost under Z (direct detection)

$$\Rightarrow \boxed{\chi = n\text{-tuplet of } SU(2)_L \quad Y = 0}$$

Stable: No renormalizable nor dim-5 operators that lead to decay

\Rightarrow first candidates are $n = 5$ fermion and $n = 7$ scalar

Lightest component neutral: $M_Q - M_{Q=0} \simeq Q(Q + \frac{2Y}{c_{\theta_w}})\Delta M$



$$\Delta M^{2\text{-loop}} = 164.5 \pm .5 \text{ MeV}$$

Ibe Matsumoto Sato 1212.5989

Minimal Dark Matter: candidates


Allowed: χ neutral under g, γ , and almost under Z (direct detection)

$$\Rightarrow \boxed{\chi = n\text{-tuplet of } SU(2)_L \quad Y = 0}$$

Stable: No renormalizable nor dim-5 operators that lead to decay

\Rightarrow first candidates are $n = 5$ fermion and $n = 7$ scalar

Lightest component neutral: $M_Q - M_{Q=0} \simeq Q(Q + \frac{2Y}{c_{\theta_w}})\Delta M$



$$\Delta M^{2\text{-loop}} = 164.5 \pm .5 \text{ MeV}$$

Ibe Matsumoto Sato 1212.5989


Avoid g_2 Landau pole before M_{Pl} \Rightarrow n not too large

In practice: $n \leq 8$ for scalars, $n \leq 5$ for fermions

[issue from 2-loop? Nardecchia et al, work in progress]



Relic abundance

Typical WIMP candidate $\rightarrow M_{\text{DM}} \sim \text{TeV}$




Important to include:

- ◊ Coannihilations
- ◊ Sommerfeld enhancement
- ? NLL corrections




γ continuum with FERMI - II


NFW profile, conservative bound




NFW profile, conservative bounds



NFW profile, including background




NFW profile, bounds including background




γ continuum with FERMI - II


Burkert profile, conservative bound




Burkert profile, conservative bounds



Burkert profile, including background



Burkert profile, bounds including background




Why an EW fermion triplet?


→ **Stable** if one imposes L or $B - L$ or discrete subgroup (already in the SM!)
[also kills all higher-dimensional operators that could make it decay]

→ Stabilizes Standard Model vacuum

without MDM [Buttazzo et al 1307.3536]



with MDM [Chao et al 1210.0491]




- Not big contribution to m_h ⇒ does not worsen fine-tuning
→ Helps with unification of gauge couplings

Why an EW fermion triplet?

→ Connection with SUSY with heavy scalars

James Wells hep-ph/0306127



Keep all good features of Supersymmetry
DM, unification of gauge couplings,...

And accept a tuned m_h (e.g. anthropic)

- All other scalars are heavier
- Higgsinos also heavier if $\mu \sim m_{3/2}$
- Wino LSP candidate for Dark Matter!

See also:

Arkani-Hamed Dimopoulos hep-th/0405159

Giudice Romanino hep-ph/0406088

...

Arvanitaki Craig Dimopoulos Villadoro 1210.0555

...

D'Eramo Hall Pappadopulo 1409.5123

Back up Extra Higgses

Extrapolation of direct searches I

We started from (and improved)

- i) Collider Reach (β) Salam Weiler 2014
- ii) Thamm Torre Wulzer 1502.01701

m_0 excluded at LHC8, obtain m_1 at future collider via

$$B(s_1, L_1, m_1) = B(s_0, L_0, m_0)$$

$$B(s, L, m) \propto L \times \int d\hat{s} \frac{1}{\hat{s}} \hat{s} \hat{\sigma}(\hat{s}) \frac{d\mathcal{L}}{d\hat{s}}(s)$$

Extrapolation of direct searches I

We started from (and improved)

- i) Collider Reach (β) Salam Weiler 2014
- ii) Thamm Torre Wulzer 1502.01701

m_0 excluded at LHC8, obtain m_1 at future collider via

$$B(s_1, L_1, m_1) = B(s_0, L_0, m_0)$$

$$B(s, L, m) \propto L \times \frac{\Delta \hat{s}}{m^2} \textcolor{red}{c} \left. \frac{d\mathcal{L}}{d\hat{s}}(s) \right|_{\hat{s}=m^2}$$

$\hat{s}\hat{\sigma}(\hat{s}) = c \Rightarrow \frac{d\mathcal{L}}{d\hat{s}}$ drives the reach

$$L_1 \textcolor{red}{c}_{ij} \left. \frac{d\mathcal{L}_{ij}}{d\hat{s}}(s_1) \right|_{\hat{s}=m_1^2} = L_0 \textcolor{red}{c}_{ij} \left. \frac{d\mathcal{L}_{ij}}{d\hat{s}}(s_0) \right|_{\hat{s}=m_0^2}$$

Extrapolation of direct searches I



We started from (and improved)

- i) Collider Reach (β) Salam Weiler 2014 ii) Thamm Torre Wulzer 1502.01701

m_0 excluded at LHC8, obtain m_1 at future collider via $B(s_1, L_1, m_1) = B(s_0, L_0, m_0)$

$$B(s, L, m) \propto L \times \frac{\Delta \hat{s}}{m^2} \textcolor{red}{c} \left. \frac{d\mathcal{L}}{d\hat{s}}(s) \right|_{\hat{s}=m^2} \quad \hat{s}\hat{\sigma}(\hat{s}) = \textcolor{red}{c} \Rightarrow \frac{d\mathcal{L}}{d\hat{s}} \text{ drives the reach}$$

$$L_1 \textcolor{red}{c}_{ij} \left. \frac{d\mathcal{L}_{ij}}{d\hat{s}}(s_1) \right|_{\hat{s}=m_1^2} = L_0 \textcolor{red}{c}_{ij} \left. \frac{d\mathcal{L}_{ij}}{d\hat{s}}(s_0) \right|_{\hat{s}=m_0^2}$$



Extrapolation of direct searches I

We started from (and improved)

- i) Collider Reach (β) Salam Weiler 2014 ii) Thamm Torre Wulzer 1502.01701



m_0 excluded at LHC8, obtain m_1 at future collider via

$$B(s_1, L_1, m_1) = B(s_0, L_0, m_0)$$



$$B(s, L, m) \propto L \times \frac{\Delta \hat{s}}{m^2} \textcolor{red}{c} \left. \frac{d\mathcal{L}}{d\hat{s}}(s) \right|_{\hat{s}=m^2}$$

$\hat{s}\hat{\sigma}(\hat{s}) = \textcolor{red}{c} \Rightarrow \frac{d\mathcal{L}}{d\hat{s}}$ drives the reach

$$L_1 \textcolor{red}{c}_{ij} \left. \frac{d\mathcal{L}_{ij}}{d\hat{s}}(s_1) \right|_{\hat{s}=m_1^2} = L_0 \textcolor{red}{c}_{ij} \left. \frac{d\mathcal{L}_{ij}}{d\hat{s}}(s_0) \right|_{\hat{s}=m_0^2}$$





Extrapolation of direct searches II



Assumptions/limitations

- Not valid if systematics dominate and change significantly from s_0 to s_1

Extrapolation of direct searches II



Assumptions/limitations

- Not valid if systematics dominate and change significantly from s_0 to s_1
- $\hat{s} \gg m_{\text{bkg}}$ [i.e. not valid at $\hat{s} \sim 2m_t$ for $\phi \rightarrow hh(4b)$]
- $\frac{\Delta\hat{s}}{m^2} \ll 1$ i.e. not valid if analysis depends a lot on shape far from peak


An extra doublet-like state H

$$[\gamma, \sigma = 0, \quad m_{h_2} \gg m_{h_1}, m_{h_3}]$$


Barbieri Buttazzo Kannike Sala Tesi 1304.3670, 1307.4937

$$\frac{g_{h_3 tt}}{g_{htt}^{\text{SM}}} = s_\delta - \frac{c_\delta}{t_\beta} \quad \frac{g_{h_3 bb}}{g_{hbb}^{\text{SM}}} = s_\delta + t_\beta c_\delta \quad \frac{g_{h_3 VV}}{g_{hVV}^{\text{SM}}} = s_\delta \quad [\Delta_t = 75 \text{ GeV}]$$

Status fit LHC8:



$m_{H^\pm} > 480 \text{ GeV}$ from $B \rightarrow X_s \gamma$!



$$[\widetilde{\mathcal{M}}_{12}^2(t_\beta, \dots) = 0 \rightarrow \delta = 0]$$

h_3 phenomenology: more similar to MSSM

see e.g. Craig et al. 1504.04630

An extra doublet-like state H


$$[\gamma, \sigma = 0, \quad m_{h_2} \gg m_{h_1}, m_{h_3}]$$

Barbieri Buttazzo Kannike Sala Tesi 1304.3670, 1307.4937


$$\frac{g_{h_3 tt}}{g_{htt}^{\text{SM}}} = s_\delta - \frac{c_\delta}{t_\beta} \quad \frac{g_{h_3 bb}}{g_{hbb}^{\text{SM}}} = s_\delta + t_\beta c_\delta \quad \frac{g_{h_3 VV}}{g_{hVV}^{\text{SM}}} = s_\delta \quad [\Delta_t = 75 \text{ GeV}]$$

Projections fit LHC14 (300 fb⁻¹):

dashed: m_{H^\pm} cont: λ





$m_{H^\pm} > 480$ GeV from $B \rightarrow X_s \gamma$!



$$[\widetilde{\mathcal{M}}_{12}^2(t_\beta, \dots) = 0 \rightarrow \delta = 0]$$

h_3 phenomenology: more similar to MSSM

see e.g. Craig et al. 1504.04630

Status fit LHC8:[dashed: m_{H^\pm} cont: Δ_t]

Red regions excluded by direct searches at LEP and CMS

Projections fit LHC14: above regions completely excluded

[if $\frac{\mu A_t}{m_t^2}$ very large, conclusions could change...]